

# MAKI SPECIMEN LABEL

## PRECAUTIONARY STATEMENTS

### HAZARD TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Keep away from humans, domestic animals and pets. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of the blood and cause bleeding.

Do not get in eyes, on skin or on clothing. Wash arms, hands and face with soap and water after applying and before eating and smoking.

#### IF SWALLOWED CALL A PHYSICIAN AT ONCE.

**NOTE TO PHYSICIAN:** If ingested, administer Vitamin K<sub>1</sub> intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

#### ENVIRONMENTAL HAZARDS

This product is toxic to mammals and birds. Do not apply this product directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

## STORAGE AND DISPOSAL

### DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.

**STORAGE:** Store in original container in a cool, dry place inaccessible to children and pets.

#### PESTICIDE DISPOSAL:

**Commercial use:** Wastes resulting from the use of this product should be securely wrapped in several layers of newspaper and then disposed of onsite or at an approved waste disposal facility.

**Household use:** Securely wrap wastes resulting from the use of this product in several layers of newspaper and discard in trash.


#### CONTAINER DISPOSAL:

**Commercial use:** Dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Household use:** Do not reuse empty container. Wrap container securely in several layers of newspaper and discard in trash.

### KEEP AWAY FROM FOODSTUFFS

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



**NORWAY RATS AND HOUSE MICE CAN CONSUME A LETHAL DOSE IN ONE FEEDING WITH FIRST DEAD RODENTS APPEARING FOUR OR FIVE DAYS AFTER TREATMENT BEGINS.**

**KILLS NORWAY RATS, ROOF RATS, HOUSE MICE AND WARFARIN-RESISTANT NORWAY RATS**

#### Active Ingredient:

*3-[3-(4'-bromo[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one	0.005%
Inert Ingredients	99.995%
Total:	100.000%

\*Bromadiolone-Supercad Patented in U.S.A.

## KEEP OUT OF REACH OF CHILDREN CAUTION

May be harmful or fatal if swallowed or absorbed through the skin. Do not get in eyes, on skin or on clothing. Wash arms, hands and face with soap and water after applying and before eating and smoking. **If swallowed, call a physician at once.** See left panel for additional precautionary statements.

MANUFACTURED BY:

**LIPHA<sup>®</sup>TECH**

LiphaTech Inc.  
3101 West Custer Ave.  
Milwaukee, WI 53209

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**IMPORTANT:** Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled, and unconsumed bait as specified on this label.

**USE RESTRICTIONS:** For control of Norway rats, roof rats, and house mice.

**URBAN AREAS:** This product may be used in and around the periphery of homes, industrial, commercial and public buildings. This product may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. This product may also be used in alleys. Do not use in sewers.

**NON-URBAN AREAS:** This product may be used inside homes and agricultural buildings.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

#### APPLICATION DIRECTIONS:

**RATS:** Apply 3 to 16 one-ounce MAKI MINI BLOCKS at each placement location. Space bait placements 15 to 30 feet apart in infested areas. Maintain an uninterrupted supply of fresh bait for 10 days or until fresh signs of rat activity cease to appear.

**FOR BURROW** applications: Insert 1 to 4 MAKI MINI BLOCKS deep into each active burrow. Check treated burrows each morning and evening to make sure that baits have not been pushed out of burrows.

**MICE:** Apply 1 one-ounce MAKI MINI BLOCK per typical placement. Two MAKI MINI BLOCKS may be needed at points of extremely high mouse activity. Space placements at intervals of 8-12 feet. Maintain an uninterrupted supply of fresh bait for 15 days or until fresh signs of mouse activity cease to appear.

**RATS/MICE:** Replace contaminated or spoiled bait immediately. Collect and properly dispose of all dead animals and leftover bait, and any bait that has become accessible to nontarget animals. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.