

RIPTIDE™

Waterbased Pyrethrin ULV

- Kills Stored Product Pests
- Kills Cockroaches
- Kills Flying Insects
- Kills Crawling Insects
- Kills Mosquitoes and Biting Flies

ACTIVE INGREDIENTS:

Pyrethrins	5.00%
Piperonyl butoxide, Technical*.....	25.00%
OTHER INGREDIENTS.....	70.00%
	100.00%

*Equivalent to 20.00% (butylcarbityl) (6-propylpiperonyl) ether and 05.00% related compounds.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

-
- IF SWALLOWED:**
- Call poison control center or doctor immediately for treatment advice.
 - Have a person sip a glass of water if able to swallow.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give anything by mouth to an unconscious person.
- IF ON SKIN OR CLOTHING:**
- Take off contaminated clothing.
 - Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.
-

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin. Wash thoroughly with soap and water after handling. Avoid contact with skin, eyes or clothing. Avoid contamination of food or feedstuffs.

Remove pets, birds and cover fish aquariums before spraying. Avoid exposure to reptiles to this product.

Do not apply space spray while food processing is underway. Cover or remove all food processing

surfaces. Thoroughly wash all food processing surfaces before reuse. After spraying in bakeries, meat packing plants, food processing plants, etc., all benches, shelving, equipment, etc. where exposed food will be handled must be washed with an effective cleaning compound followed by a potable water rinse to remove all traces of contamination. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions above in those establishments which do NOT operate under Federal meat, poultry, shell egg grading and egg products inspection programs.

Do not apply directly to food, water or feed supplements. Do not contaminate milk or milk handling equipment. Wash udder and teats of dairy animals before milking.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and other aquatic invertebrates. Do not contaminate water when disposing of equipment washwaters. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

INDOOR AREAS for use in and around: [such as]:

Animal Areas and Quarters, Animal Shelters, Animal Control Centers, Apartments, Arcades, Attics, Auditoriums, Bakeries, Beverage Plants, Bars, Basements, Bathrooms, Boats, Bottling Plants, Breweries, Brewery Warehouses, Buildings, Buses, Cabins, Cabinets, Cafeterias, Campers, Canneries, Cattle Barns, Churches, City Missions, Circus Tents, Closets, Club House, Commercial Buildings, Communication Centers, Condominiums, Concession Stands, Court Houses, Crawl Spaces, Crematoriums, Dairy and Livestock Barns, Day Care Centers, Delicatessens, Department Stores, Diners, Dog Houses, Storage or Processing Areas that contain Dried Fruit Products, Drug Stores, Dumpsters, Dwellings, Egg Processing Plants, Factories, False Ceilings, Freight Containers, Fruit Packing Sheds, Funeral Parlors, Garbage Bins, Garbage Cans, Garbage Compactors, Garbage Trucks, Gas Stations, Grain Elevators, Grain Harvesting and Handling Equipment, Garages, Grain Mills, Granaries, Grocery Stores, Homes, Horse Stables and Barns, Horse Trailers, Hospices, Hospitals, Hotels, Indoor Eating Establishments, Industrial Installations, Institutional Dining Areas, Institutions, Jails, Kennels, Laboratories, Laundromats, Lavatories, Libraries, Liquor Stores, Livestock Barns, Livestock Hauling Equipment, Loafing Sheds, Locker Rooms, Marinas, Meat Packaging Plants, Milk Rooms, Mills, Mausoleums, Meat Packing Plants, Mess Halls, Mini-Storages, Mobile Canteens, Mobile Homes, Morgues, Mortuaries, Motels, Movie Theaters, Museums, Mushroom Processing, Nursing Homes, Office Buildings, Other Public Buildings, Parking Garages, Passenger Railroad Cars, Peanut Warehouses, Pool Halls, Poultry Houses, Poultry Plants, Pounds, Prisons, Public Storage, Rabbit Processing Plants, Railroad Cars, Recreational Vehicles, Research Animal Quarters, Restaurants, Retail Stores, Rice Mills, Schools, Ships, Ship Holds, Smoke Shops, Sports Stadiums, Stables, Storage Bins (areas), Storage Areas for Food in Multi-walled Paper or Cloth Bags, Stores, Supermarkets, Swine Houses, Taverns, Textile Mills and Warehouses, Theaters, Tobacco Factories, Tobacco Warehouses, Tool Sheds, Trains, Transportation Equipment, Trucks, Truck Trailers, Tunnels, Utilities, Utility Rooms, Vending Machines, Veterinary Hospitals, Wall and Equipment Voids, Warehouses, Washrooms, Wholesale Stores, Wine Cellars, Wineries, Zoos and other similar areas.

OUTDOOR AREAS for use in and around: [such as]:

Alleys, Amphitheaters, Amusement Parks, Athletic Fields, (baseball, football, soccer), Back Yards, Balconies, Baseball Fields, Beaches, Bleachers, Bridges, Carwashes, Campgrounds, Circuses, Construction Sites, Corrals, Courtyards, Curbs, Drive-In Restaurants, Drive-In Theaters, Dumpsters, Fair Grounds, Feedlots, Front Yards, Game Fields, Garbage Dumps, Gazebos, Golf Courses, Grandstands, Junk Yards, Game Fields, Kennels, Levees, Manure Piles, Marinas, Outdoor Eating Establishments, Parks, Parking Lots, Patios, Playgrounds, Ranges, Recreational Areas, Rest Areas, Roadsides, Rooftops, Sanitary Sewers (Manholes), Side Yards, Side Walks, Stadiums, Tennis Courts, Truck Stops, Underpasses, Utility Room Outdoor, Weigh Stations, Wild Animal Parks, Wood Piles, Zoos and other similar areas.

FOR USE ON: [such as]:

Beefalo, Birds, Buffalo, Calves, Cattle [beef] [milking], Chickens, Donkeys, Ducks, Emus, Exotics such as European Red Deer, Geese, Horses, Lambs, Llamas, Mules, Ostrich, Oxen, Pheasants, Pigs, Ponies, Pullets, Quail, Ratites, Roosters, Sheep, Swine, Turkeys.

FOR THE CONTROL OF INSECTS including: [such as]:

Almond Moths, Angoumois Grain Moths, Ants, Black Fly, Blood Sucking Lice, Boxelder Bugs, Brown Dog Ticks, Cadelles, Carpet Beetles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Clothes Moths, Clover Mites, Cockroaches, Coffee Bean Weevils, Confused Flour Beetles, Crickets, Dark Mealworms, Darkling Beetles, Deer Flies, Deer Ticks, Dermestid Beetles, Dried Fruit Beetle, Drugstore Beetles, Earwigs, Face Fly, Firebrat, Flat Grain Beetles, Fleas, Flies, Fruit Flies, Gnats, Grain Mites, Granary Weevils, Horn Flies, Hornets, Horseflies, Indian Meal Moths, Khapra Beetles, Lesser Grain Borers, Lesser House Fly, Mediterranean Flour Moths, Merchant Grain Beetle, Midges, Millers, Millipedes, Mosquitoes, Flying Moths, Palmetto Bugs, Pillbugs, Red Flour Beetles, Rice Flour Beetles, Rusty Grain Beetles, Saw-toothed Grain Beetle, Silverfish, Skipper Fly, Spiders, Spider Beetles, Stable Flies, Sowbugs, Ticks, Ticks that may carry and transmit Lyme Disease, Tobacco Moths, Trogoderma Beetles, Warehouse Beetles, Wasps, Waterbugs, Yellow Jackets, Yellow Mealworms.

INDOOR USE—UNDILUTED

ULV AND SPACE SPRAY: Use in aerosol generators to kill exposed accessible stages of: Flies (including Fruit Flies), Mosquitoes, Small Flying Moths, Fleas, Gnats, Wasps, Hornets, Clover Mites, Cheese Mites, Cheese Skippers, Boxelder Bugs, Earwigs, Ants and the following common Stored Product Pests: Granary Weevils, Rice Weevils, Confused Flour Beetles, Saw-toothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths, Tobacco Moths, Yellow Meal Worms, Dark Meal Worms, Grain Mites, Cadelles and Red Flour Beetles. Close room and shut off all air conditioning or ventilating equipment. Apply at the rate of 1/2 to 1 fluid ounce concentrate per 1,000 cubic feet of space. Direct spray toward upper corners of room and ceiling. Leave room closed for 1/2 hour and ventilate thoroughly before re-entry.

CONTACT AND SPACE SPRAY: To kill Cockroaches, Silverfish, Earwigs, Ants and Spiders: use a mechanical aerosol generator or a fogger adjusted to deliver aerosol droplets. Direct the spray first into all crack and crevices, behind all equipment, cupboards, obstructions or dark harborage areas, walls, floors, underneath sinks, behind pipes and in all places that harbor insects to provide maximum penetration and impingement on the insects. Contact as many insects as possible. Do not allow spray to contact stored food. Then treat as a space spray, directing the spray into upper corner areas of room and ceiling, apply at a rate of 1 fluid ounce per 1,000 cubic feet of space. Facilities and equipment in federally inspected meat and poultry plants must be washed with an effective cleaning compound and rinsed with potable water after spraying. Vacate treated area and leave the room closed for 1 hour. Ventilate thoroughly before re-entry.

DRIED FRUIT PRODUCT STORAGE AREAS: In storage or being processed for control of Almond Moths, Ants, Cereal Beetles, Chocolate Moths, Cigarette Beetles, Cockroaches, Confused Flour Beetles, Crickets, Dried Fruit Beetles, Earwigs, Flies, Fruit Flies, Fungus Gnats, Indian Meal Moths, Khapra Beetles, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles, Silverfish and Warehouse Beetles. Do not apply to fruit directly. Apply as a fine mist or fog in the air above and around the trays, bins and shelves at the rate of 1 fluid ounce per 1,000 cubic feet of space. Repeat application at this dosage as required.

TREATMENT OF STORED FOOD AREAS: Foods in multi-walled bags, cloth bags or other packaging stored in such areas, listed areas for the control of accessible adult stages of: Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Coffee Bean Weevils, Confused Flour Beetles, Crickets, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Earwigs, Firebrats, Flies, Fruit Flies, Fungus Gnats, Grain Mites, Granary Weevils, Hide Beetles, Indian Meal Moths, Khapra Beetle, Lesser Grain Beetles, Mediterranean Flour Moths, Mosquitoes, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Saw-toothed Grain Beetles, Silverfish, Small Flying Moths, Spiders, Spider Beetles, Warehouse Beetles and Yellow Mealworms. Infested stored products should be destroyed, fumigated or treated by other effective methods using an approved product intended for this purpose. As noted above, disperse first by directing spray into cracks and crevices, corners, pallets and around stacks of packaged goods hitting as many insects as possible and then apply as a space spray at a dosage of 1 fluid ounce per 1,000 cubic feet of room space. Do not remain in treated area and ventilate before re-entry.

ON PEANUTS, NUTS (ALMONDS AND WALNUTS), BEANS, (INCLUDING COCOA), SEEDS (BIRDSEED MIXTURES, COTTONSEED AND FLAXSEED) AND COPRA IN BULK, BAGS OR OTHER PACKAGING: For the control of accessible stages of: Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Coffee Bean Weevils, Confused Flour Beetles, Crickets, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Earwigs, Firebrats, Flies, Fruit Flies, Fungus Gnats, Grain Mites, Granary Weevils, Hide Beetles, Indian Meal Moths, Khapra Beetles, Lesser Grain Beetles, Mediterranean Flour Moths, Mosquitoes, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Saw-toothed Grain Beetles, Silverfish, Small Flying Moths, Spiders, Spider Beetles, Warehouse Beetles, Yellow Mealworms and other listed insects apply to the surface of stored or bagged products at the rate of 1/4 fluid ounce per 1,000 square feet. Also spray walls, floors and other surfaces of bins, storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Then apply as a space spray by directing the spray toward the ceiling and upper corners of the area and behind any obstructions. Apply at the rate of 1 fluid ounce per 1,000 cubic feet of room space. Do not remain in treated area and ventilate before re-entry.

INDOOR USE—DILUTED

SPACE SPRAY: For use in mechanical fogging or spraying equipment to control accessible stages of Almond Moths, Angoumois Grain Moths, Ants, Cadelles, Cereal Beetles, Cheese Mites, Cheese Skippers, Chocolate Moths, Cigarette Beetles, Cockroaches, Coffee Bean Weevils, Confused Flour Beetles, Crickets, Dark Mealworms, Dried Fruit Beetles, Drugstore Beetles, Earwigs, Firebrats, Flies, Fruit Flies, Fungus Gnats, Grain Mites, Granary Weevils, Hide Beetles, Indian Meal Moths, Khapra Beetles, Lesser Grain Beetles, Mediterranean Flour Moths, Mosquitoes, Red Flour Beetles, Rice Weevils, Rusty Grain Beetles, Saw-toothed Grain Beetles, Silverfish, Small Flying Moths, Spiders, Spider Beetles, Warehouse Beetles and Yellow Mealworms. Remove or cover exposed food and cover food handling surfaces in edible product areas of food handling establishments. Close room and shut off all air conditioning or ventilating equipment. Dilute at the rate of 1 part concentrate into 1 to 49 parts water and mix well. Apply at the rate of 1-2 fluid ounce per 1,000 cubic feet of space, filling the room with mist. Vacate treated area and ventilate before re-entry.

FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS: Dilute one part concentrate with 4 parts water and use with a ULV spraying device at the rate of one fluid ounce per 1,000 cubic feet of room space. Apply the spray into all cracks and crevices in woodwork,

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ALL DIRECTIONS COMPLETELY BEFORE USE AND FOLLOW ALL DIRECTIONS AND PRECAUTIONS WHEN USING PRODUCT.

For best results, follow directions for specific use areas.

walls, floors, underneath sinks, behind pipes and in all places that shelter insects. Contact as many insects as possible. Before use, exposed food products and packaging materials must be covered or removed. After use, equipment and utensils must be washed with an effective cleaning compound and rinsed with potable water. Use as a space spray only when plant is not in operation. Repeat as necessary but not more than 1 time per day. Do not reapply within 1 day.

SURFACE SPRAY: To control accessible stages of: Cockroaches, Silverfish, Beetles and Crickets: Dilute 1 part concentrate into 1 to 49 parts water, mix well and apply as a coarse wetting spray using approximately 1 gallon of spray mixture per 750 square feet of surface. To ensure maximum control of listed crawling insects, spray thoroughly into all cracks, molding, crevices, walls, floors and other surfaces of bins, storage and handling areas. Particular attention should be paid to areas where moisture collects such as drains, sinks and water pipes, around the base of heavy machinery and equipment contacting as many insects as possible.

ANTS: Spray directly into nests, along trails and directly on all ants observed.

FOR IN TRANSIT PROTECTION: To control accessible stages of listed insects, clean and spray truck beds, boxcars, ships holds and other listed transit equipment before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers lightly as stacked. Apply diluted spray at the rate of 1 gallon per 1,000 square feet.

TREATMENT OF STORED GRAIN: To control accessible stages of: Indian Meal Moth, Angoumois Grain Moths: Apply diluted spray to the surface of binned grain at a rate of 0.5 gallons per 1,000 cubic feet. Apply at monthly intervals with the first appearance of infestation. In severe infestations of flying moths, break up webbing with a rake before spraying and make a second application in two weeks.

STORAGE AREAS FOR DRIED FRUIT PRODUCTS for control of: Confused Flour Beetles, Dried Fruit Beetles, Indian Meal Moths and Saw-toothed Grain Beetles. Remove and destroy infested products. Apply diluted spray as a fine mist in the air above the trays and shelves. Do not apply to fruit directly. Use 1 fluid ounce per 1,000 cubic feet of space. Repeat application at this dosage in any given month. Leave the room closed for one hour. Do not remain in treated areas and ventilate the area before re-entry. Sweep up and destroy fallen insects.

FOOD STORAGE AREAS containing multi-walled bags, cloth bags and other packaging in warehouses, storage rooms, flour mills, grain elevators, granaries and other listed areas for the control of: Angoumois Grain Moths, Cadelles, Cheese Mites, Cheese Skippers, Cigarette Beetles, Confused Flour Beetles, Dark Meal Worms, Grain Mites, Granary Weevils, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Rice Weevils, Saw-toothed Grain Beetles and Yellow Meal Worms. Remove and destroy infested products. Disperse diluted spray first by directing spray into corners, under pallets and around stacks of packaged foods hitting as many insects as possible and then treat as a space spray. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose. Use at a rate of 1-2 fluid ounces per 1,000 cubic feet of space. Do not remain in treated areas and ventilate area before re-entry. This application will kill exposed stages of those stored product pests listed. Sweep up and destroy fallen insects.

FOR USE ON ANIMALS: To protect: Beef and dairy cattle, horses from Horn Flies, House Flies, Mosquitoes and Gnats, dilute 1 part concentrate with 49 parts water, mix well and apply a light mist sufficient to wet the tips of the hair. To control Stable Flies, Horse Flies and Deer Flies on animals apply 2 fluid ounces per adult animal, sufficient to wet the hair but not to soak the hide. Repeat treatment once or twice daily or at intervals to give continued protection. To control the motile stages of Blood Sucking Lice on beef and dairy cattle and other listed animals, apply the spray to thoroughly wet the hair. Use 2 fluid ounces or less per animal and repeat every 2 to 3 weeks as required.

To control Poultry Lice, spray roosts, walls and nests or cages thoroughly with a dilution of 1 part concentrate with 25 parts water. This should be followed by spraying over the birds with a fine mist.

ANIMAL QUARTER USE - DAIRY AND BEEF BARNs, POULTRY HOUSES, SWINE HOUSING AND OTHER LISTED AREAS: For use in and around listed use areas to control accessible stages of: Brown Dog Ticks, Clover Mites, Cockroaches, Crickets, Darkling Beetles, Deer Flies, Earwigs, Fannia Flies, Face Flies, Firebrats, Fleas, Flies, Horn Flies, Hornets, Mosquitoes, Silverfish, Small Flying Moths, Spiders, Stable Flies, Ticks, Wasps and Yellow Jackets.

As a space spray in conventional ULV cold aerosol generator, automatic ULV misting systems and equipment, or fogger adjusted to deliver aerosol size droplets. Whenever possible, space spray application should be made in the late evening when insects are at rest. Apply undiluted at the rate of 0.25-1.0 fluid ounces per 1,000 cubic feet of space above the animals. This concentrate may also be diluted with water at the rate of 1 part concentrate to 10 parts water and applied at the rate of 1-2 fluid ounces per 1,000 cubic feet of space above the animals. Direct spray toward the upper portions of the enclosure above the animals, filling the room with mist or fog. Vacate treated area and ventilate before re-occupying.

To control Bedbugs and Mites in poultry houses, prepare a dilution of 1 part concentrate to 25 parts water and spray crevices of roost poles, cracks in walls and cracks in nests where the Bedbugs and Mites hide.

OUTDOOR USE—UNDILUTED

To control accessible stages of listed flying insects in listed outdoor areas. If possible, space spray application should be made in the late evening when insects are at rest. With a ULV spraying device, spray at a dosage rate of 5 fluid ounces per minute.

TEMPORARY REDUCTION OF ANNOYANCE from: Flies, Mosquitoes and Small Flying Moths. The afore-mentioned application for outdoor ground application will afford temporary reduction of annoyance from these pests in listed outdoor areas. Direct application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage. In areas such as zoos, cover water, drinking fountains and animal feed before use.

OUTDOOR USE—DILUTED

HOSE-END APPLICATIONS - TO CONTROL PESTS ON SHRUBS, FRUIT AND SHADE TREES:

Pests: Aphids, Apple and Thorn Skeletonizers, Bagworms, Bean Beetles, Cabbageworms, Codling Moths, Cutworms, European Pine Shoot Moths, Flea Beetles, Hemlock Wolly Adelgids, Holly Bud Moths, Lacebugs, Leafhoppers, Leafminers, Leafrollers, Mealybugs, Oak Loopers, Obscure Root Weevils, Omnivorous Leaf-tiers, Pear Slugs, Scale Crawlers, Spider Mites, Tent Caterpillars, Thrips, Webworms and Whiteflies.

Plants: Birch, Crabapple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Hemlock, Holly, Juniper, Lilac, Locust, Maple, Oak, Plum, Pine, Poplar, Spruce, Sycamore, Willow and Yew.

Mixing Instructions: For Shrubs and Shade Trees follow directions on hose end applicator to obtain 4 tbsp (2 fl. oz.) per gallon of water.

1 Tablespoon (tbsp) = 3 teaspoons (tsp)

1 fluid ounce = 2 tbsp

How To Apply: Spray plant to wet the leaves and branches until dripping.

AROUND HOUSE FOUNDATION:

Nuisance Pests Outdoors: Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Scorpions, Sowbugs, Spiders and Springtails.

Mixing Instructions: For Around House Foundation follow directions on hose end applicator to obtain 4 tbsp (2 fl. oz.) per gallon of water.

1 Tablespoon (tbsp) = 3 teaspoons (tsp)

1 fluid ounce = 2 tbsp

How To Apply: To keep insects from entering the house, spray a 5 feet band of soil around the house next to the foundation. Also spray the foundation of the house to a height of 2 feet. Also spray walks, firewood piles, garbage cans, tree trunks and into cracks and other places where insects can hide.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry place. Keep container closed.

DISPOSAL:

If empty: Do not reuse container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.