



- **A micro-emulsion, water-based spray**
- **Kills Mosquitoes, including the Culex species that may carry and transmit the West Nile Virus**
- **Kills and repels Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Mosquitoes and Gnats**
- **Controls Deer Ticks and other Ixodid species that may carry and transmit Lyme Disease**
- **Vampyre™ Misting Concentrate provides rapid kill of insects without objectionable residues**

ACTIVE INGREDIENT:

Pyrethrins	3.00%
*Piperonyl butoxide, Technical	30.00%
OTHER INGREDIENTS	67.00%
	100.00%

*Equivalent to min. 24% (butylcarbityl) (6-propylpiperonyl) ether and 6% related compounds

KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside for first aid and precautionary statements.

For the control of insects, including, but not limited to:

Almond Moths	Cigarette Beetles	Flat Grain Beetles	Lice	Saw-toothed Grain Beetles
Angoumois Grain Moths	Clothes Moths	Fleas	Mediterranean Flour Moths	Sheep Ticks
Ants	Clover Mites	Flies	Merchant Grain Beetles	Silverfish
Bedbugs	Cockroaches	Fruit Flies	Midges	Skipper Flies
Black Carpenter Ants	Confused Flour Beetles	Fungus Gnats	Miller Moths	Small Flying Moths
Black Flies	Crickets	Gnats	Millipedes	Sowbugs
Black Widow Spiders	Dark Mealworms	Grain Mites	Mosquitoes including the Culex species that may carry and transmit the West Nile Virus	Spider Beetles
Blood Sucking Lice	Darkling Beetles	Granary Weevils	Hide Beetles	Spiders
Boxelder Bugs	Deer Flies	Horse Flies	Horn Flies	Stable Flies
Brown Dog Ticks	Deer Ticks	Hornets	Horse Flies	Ticks that may carry and transmit Lyme Disease
Cadelles	Dermestid Beetles	House Flies	Indian Meal Moths	Tobacco Moths
Carpet Beetles	Dog Fleas	Lesser Grain Beetles	Khapra Beetles	Trogoderma Beetles
Cat Fleas	Dried Fruit Beetles	Lesser Grain Borers	Lesser Grain Beetles	Warehouse Beetles
Cellar Spiders	Drugstore Beetles	Lesser House Flies	Lesser House Flies	Wasps
Cereal Beetles	Earwigs	Face Flies	Face Flies	Waterbugs
Cheese Mites	Fannia Flies	Chocolate Moths	Chocolate Moths	Yellow Mealworms
Cheese Skippers	Firebrat			Yellow-Jackets

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read all directions completely before use. For best results, follow directions for specific use areas.

GENERAL:

This concentrated product must be diluted with water. When making dilutions, add concentrate to the tank and add water. Agitate well while adding water and apply as directed.

Confine use of this product in food processing establishments to time periods when the plant is not in operation. Remove or cover food during treatment. Cover all food processing surfaces during treatment or thoroughly clean before use following treatment.

For maximum effectiveness, following the dilution chart and using a combination of spot, surface, and fogging treatment are recommended.

0.05% Pyrethrins	add 1 part concentrate to 59 parts water or 2 fluid ounces concentrate / gallon water
0.10% Pyrethrins	add 1 part concentrate to 29 parts water or 4.3 fluid ounces concentrate / gallon water
0.30% Pyrethrins	add 1 part concentrate to 9 parts water or 13 fluid ounces concentrate / gallon water
0.50% Pyrethrins	add 1 part concentrate to 5 parts water or 22 fluid ounces concentrate / gallon water

INDOOR AREAS for use in but not limited to:

Animal Areas	Commercial Buildings	Goat Holding Areas	Mess Halls	Schools
Animal Control Centers	Condominiums	Grain Elevators	Milk Houses	Seed Storage
Animal Quarters	Crawl Spaces	Grain Handling Equipment	Milk Rooms	Sheep Holding Areas
Animal Shelters	Credenzas	Grain Harvesting Equipment	Milk Sheds	Ships
Apartments	Crematoriums	Grain Mills	Milking Rooms	Ships Holds
Atriums	Dairies	Grain Storage	Mini Storage	Ships Holds
Attics	Dairy Livestock Barns	Granaries	Mobile	Smoke Shops
Auditoriums	Day Care Centers	Grocery Stores	Canteens	Solariums
Automobiles	Decks	Hog Barns	Mobile Homes	Sport Stadiums
Bakeries	Delicatessens	Homes	Morgues	Stable Bins
Bars	Dens	Horse Stables	Mortuaries	Stables
Basements	Department Stores	Horse Stables & Barns	Motels	Storage Areas
Bathrooms	Diners	Horse Trailers	Movie Theaters	Stored Food (bags)
Bedding	Dog Houses	Hospices	Museums	Stored Products
Bedrooms	Dog Sleeping Quarters	Hospitals	Nursing Homes	Stores
Beef Barns	Dormitories	Hospitals	Office Buildings	Sun Porches
Beverage Plants	Draperies	Hotels	Other Public Buildings	Sun Rooms
Boats	Dressers	Household Contents	Outdoor Eating Buildings	Swine Houses
Bookcases	Dried Fruits in Storage	Houses	Pantries	Taverns
Bottling Plants	Drive-Ins	Indoor Eating Buildings	Parking Garages	Tents
Breweries	Drugstores	Industrial Installations	Pet Bedding	Textile Mills
Brewery Warehouses	Dumpsters	Industrial Plants	Pet Carriers	Theaters
Buildings	Dwellings	Institution Dining Areas	Pet Grooming Parlors	Tobacco Factories
Housing Animals	Egg Processing Plants	Institutions	Pet Sleeping Areas	Tobacco Warehouses
Buildings Housing Poultry	Fabric Storage Areas	Jails	Pet Stores	Tool Sheds
Buses	Factories	Kennels	Planes	Trailers
Cabinets	False Ceilings	Kitchens	Play Houses	Trains
Cabins	Farm Structures	Laboratories	Play Rooms	Transportation Vehicles
Cafeterias	Floors	Lanais	Pool Halls	Tree Houses
Calving Barns	Food Processing Plants	Laundromats	Porches	Tree Nuts Storage Area
Camp Sites	Food Storage Areas	Lavatories	Poulties	Truck Trailers
Campers	Freight Containers	Libraries	Poultry Plants	Trucks
Canneries	Fruit Packing Sheds	Liquor Stores	Pounds	Tunnels
Carpets	Furniture	Livestock Barns	Prisons	Utilities
Cat Sleeping Quarters	Garages	Livestock Hauling Equipment	Public Storage	Utility Rooms
Cattle Barns	Garbage Bins	Livestock Holding Areas	Quonset Huts	Vending Machines
Chests	Garbage Cans	Living Rooms	Rabbit Processing Plants	Verandas
Churches	Garbage Compactors	Loafing Shed	Railroad Cars	Veterinary Clinics
Circus Tents	Gas Stations	Locker Rooms	Recreational Vehicles	Voids in Attics
City Missions	Gazebos	Marinas	Research	Warehouses
Closed Porches		Mattresses	Animal Quarters	Washrooms
Closets		Mausoleums	Restaurants	Wheat Mills
Clothes Storage		Meat Packing Plants	Retail Stores	Wholesale Stores
Club Houses			Rice Mills	Wine Cellars
			Rooms	Wineries
			Rugs	Zoos

OUTDOOR AREAS for uses in and around, but not limited to:

Alleys	Bridges	Fair Grounds	Outdoor Eating Establishments	Side Walks
Amphitheaters	Campgrounds	Feedlots	Parking Lots	Side Yards
Amusement Parks	Carwashes	Front Yards	Parks	Stadiums
Ant Hills	Circuses	Game Fields	Patios	Tennis Courts
Back Yards	Construction Sites	Garbage Dumps	Playgrounds	Truck Stops
Balconies	Corrals	Gazebos	Ranges	Underpasses
Baseball Fields	Courtyards	Grandstands	Recreational Areas	Utility Rooms
Beaches	Curbs	Junk Yards	Rest Areas	Outdoors
	Drive-In Restaurants	Kennels	Roadsides	Weigh Stations
	Drive-In Theaters	Levees	Rooftops	Wild Animal Parks
	Dumpsters	Manholes	Sanitary Sewers	Wood Piles
		Manure Piles		Zoos
		Marinas		

FOR USE ON but not limited to:

Beefalo	Cattle-Beef	Exotics such as European Red Deer	Mules	Quail
Birds	Cattle-Dairy	Geese	Ostrich	Ratites
Buffalo	Chickens	Horses	Oxen	Roosters
Calves	Donkeys	Lambs	Pheasants	Sheep
	Ducks	Llamas	Pigs	Swine
	Emus		Pullets	Turkeys

DIRECTIONS FOR USE

SPACE SPRAYING: Use with conventional mechanical or compressed air equipment to kill exposed accessible stages of listed insects. Close room and shut off air conditioning or ventilating equipment. Apply at the rate of 1 to 10 fluid ounces per 1000 cubic feet of space and direct spray toward upper corners of room and ceiling filling the room with mist. Leave room closed for 1/2 hour and ventilate thoroughly before re-entry.

SURFACE SPRAYING: To kill Cockroaches, Black Carpenter Ants, Silverfish, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Saw-toothed Grain Beetles, Spider Beetles, Cigarette Beetles and other listed insects, use a sprayer adjusted to a coarse wet spray. Mix well and apply as a coarse wet spray using approximately 1 gallon of spray mixture per 750 square feet of surface. Direct the spray first into all cracks and crevices, behind all equipment, cupboards, obstructions or dark harborage areas on walls and floors, underneath sinks, behind pipes, beneath stoves and refrigerators, around garbage cans and cabinets, along baseboards, door sills, window sills, door and window frames and floors, around and on drains, pipes, plumbing, behind bookcases, storage and other utility installation areas and in all places that harbor insects to provide maximum penetration and impingement on the insects. Contact as many insects as possible. Do not allow spray to contact stored food. If surface application only is to be used, spray floor and other surfaces. To kill ants, treat around doors, window frames and other areas of entry. Spray on ants whenever possible.

CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply liberally to empty chests, closets, bureaus and other storage areas, directing the spray into cracks, joints, and crevices.

CARPET BEETLES: Directly spray as many insects as possible. Spray all surfaces of upholstered furniture. Where possible, both top and bottom of rugs and carpets should be sprayed thoroughly. Also treat shelves, closets, floors, walls, windowsills, baseboards, areas under furniture, and wherever these insects are seen or suspected.

BEDBUGS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, floors, walls, furniture and other infested areas.

FOR USE IN CANNERIES: Thoroughly clean floors, conveying equipment, and walls, paying special attention to cracks and crevices, prior to bringing in produce. Follow dilution chart and spray the entire space inside the cannery including walls, ceilings, and floors, paying special attention to forcing the spray into all cracks and crevices for control of Ants, Cockroaches, Silverfish, Crickets, Spiders, Fruit Flies, Moths, House Flies, Cheese Mites and other listed insects.

FOR USE IN FOOD AND NONFOOD AREAS OF FOOD PROCESSING PLANTS, Industrial Installations, Bakeries, Rice and Wheat Mills, Restaurants, Tobacco Warehouses, Grain Elevators and Warehouses. Spray as directed to control listed flying and crawling insects. The use of this product in food processing or food handling establishments should be confined to time periods when the plant is not in operation. Food should be covered or removed during treatment.

DRIED FRUIT PRODUCTS in storage or being processed: For control of listed insects. Remove and destroy infested products. Do not apply directly to fruit. Apply as a fine mist or fog in the air above and around trays, bins, and shelves at the rate of 1 to 10 fluid ounces per 1000 cubic feet of space. Repeat application at this dosage as required. Leave the room closed for one hour. Do not remain in treated areas, and ventilate the area before re-entry. Sweep up and destroy fallen insects.

TREATMENT OF STORED FOOD AREAS: Foods in multi-walled bags, cloth bags or other packaging stored in such areas as, but not limited to listed areas. For the control of listed pests. Infested stored products should be destroyed, fumigated or treated by other effective methods using an approved product intended for this purpose. As noted above, disperse first by directing spray into cracks and crevices, corners, pallets and around stacks or packaged goods, hitting as many insects as possible. Then apply as a space spray at a dosage of 1 to 10 fluid ounces per 1000 cubic feet of room space. Repeat application as necessary for insect control. Do not remain in treated area and ventilate before re-entry. Sweep up and destroy fallen insects.

ON PEANUTS, NUTS, BEANS, SEEDS IN BULK, BAGS OR OTHER PACKAGING: For the control of listed insects. Apply to the surface of stored or bagged products. Also spray walls, floors and other surfaces of bins, storage and handling areas thoroughly, being especially careful to treat cracks, crevices and similar hiding places. Then apply as a space spray at the rate of 1 to 10 fluid ounces per 1000 cubic feet by directing the spray toward the ceiling and upper corners of the area and behind any obstructions.

FOR USE IN FEDERALLY INSPECTED MEAT AND POULTRY PLANTS: Apply with conventional mechanical or compressed air equipment at the rate of 1 to 10 fluid ounces per 1000 cubic feet of room space. Apply spray into all cracks and crevices in woodwork, walls, floors, and underneath sinks, behind pipes and in all places that shelter insects. Contact as many insects as possible. Repeat as necessary. Before use, exposed food products and packaging materials must be covered or removed. After use, equipment and utensils must be washed with an effective cleaning compound and rinsed with potable water. Use as a space spray only when plant is not in operation.

FOR IN TRANSIT PROTECTION: To control accessible stages of listed insects, clean and spray truck beds, boxcars, ship holds and other listed transit equipment before loading. Thoroughly treat walls and floors, and lightly spray each layer of stacked cartons or containers. Apply at the rate of 1 gallon per 750 square feet.

FOR USE IN LISTED PET AREAS: For best results remove, destroy or wash pet's old bedding. To reduce infestation, thoroughly spray pet (dog, cat, puppy, kitten, or animal) bedding, sleeping quarters, kennel, around baseboards, window and door frames, wall cracks, local areas of floors, and other resting places until the surface is slightly moist. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 18 inches. Also spray the entire inside surface of doghouses. Put fresh bedding in pet's quarters after spray has dried. Allow treated areas to dry thoroughly before allowing pets to re-enter.

Remove pets and birds and cover fish aquariums before spraying.

FOR KENNELS AND OTHER OUTDOOR ANIMAL AREAS: Follow dilution chart to control listed insects. Apply liberally to surfaces, paying special attention to cracks and crevices and other insect hiding places. Allow treated areas to dry thoroughly before allowing pets (animals) to re-enter.

FOR USE ON LIVESTOCK: Dilute as follows: 0.10% Pyrethrins – Add 1 part concentrate to 29 parts water or 0.50% Pyrethrins – Add 1 part Concentrate to 5 parts water and apply as directed:

- To protect beef cattle, dairy cattle, horses and other livestock from attacks and to kill and repel Stable Flies, Horse Flies, Deer Flies, House Flies, Horn Flies, Lesser House Flies, Mosquitoes including the Culex species that may carry and transmit the West Nile Virus, and Gnats, spray about 3 seconds on each side of animal, being careful to spray back, withers and forelegs thoroughly.
- To protect from Face Flies, spray the face and hair or spray on a cloth and rub on the face, being certain to avoid contact with the eyes. Repeat treatment as necessary when flies are troublesome.
- For effective control of Biting and Sucking Lice on cattle, horses, sheep, goats and hogs: spray to thoroughly wet the hair of the animal, including the head and brush of the tail. Repeat treatment in 10 days to kill newly hatched lice.
- To control Poultry Lice and exposed stages of Darkling Beetles, thoroughly spray roosts, walls, nests, and cages. Follow by spraying the birds with a fine mist.
- For control of Bedbugs and Mites on poultry and in poultry houses, spray crevices of roost poles, cracks in walls and cracks in nests where bedbugs and mites hide. Follow by spraying the birds with a fine mist.
- To control Sheep "Tick" or Ked, thoroughly wet all portions of the body by spraying directly into the wool. Treat at a rate sufficient to wet the animal.
- To kill Fleas and Ticks on livestock and to obtain protection against reinfestation, spray to sufficiently wet animal.

FOR USE IN ANIMAL QUARTERS, Barns, Dairies, Milking Parlors, Milking Rooms, Poultry Houses and other listed animal areas. For rapid control (kill) of Flies, Mosquitoes including the Culex species that may carry and transmit the West Nile Virus, Gnats and other listed flying insects: dilute with water to give a 0.10% or 0.50% concentration, and apply as a fine mist or spray. Direct spray towards the ceiling and upper corner of room until the area is filled with mist, using about 2 fluid ounces per 1000 cubic feet of space. Do not direct spray towards animals or birds. For best results, close doors and windows before spraying and keep them closed for 10 to 15 minutes following application. Applicator should vacate the treated area and ventilate it prior to reoccupying. Do not use in milk rooms where raw milk may be exposed.

FOR USE WITH HAND-HELD MECHANICAL FOGGERS to control Houseflies, Horn Flies, Stable Flies, Horse Flies, Mosquitoes including the Culex species that may carry and transmit the West Nile Virus, Gnats and other listed insects. Dilute concentrate and apply as a fine mist or fog. Direct spray toward the ceiling and upper corners of the room until the area is filled with mist, using about 2 fluid ounces per 1000 cubic feet of space. For best results, close doors and windows before spraying and keep them closed for 10 to 15 minutes after spraying. Applicator should vacate the treated area and ventilate it before reoccupying.

FOR USE IN AUTOMATIC ULV SPRAYING SYSTEMS, IN AREAS SUCH AS ANIMAL QUARTERS, DAIRY AND BEEF BARNS, POULTRY HOUSES, SWINE HOUSING, WAREHOUSES, RESIDENTIAL BACKYARDS AND OTHER LISTED INDOOR AND OUTDOOR AREAS.

Use to control, or temporarily reduce annoyance from accessible stages of: Mosquitoes including the Culex species that may carry and transmit the West Nile Virus, Flies, Gnats, Face Flies, Fannia Flies, Deer Flies, Horse Flies, Stable Flies, Horn Flies, Hornets, Wasps, Yellow Jackets, Firebrats, Small Flying Moths, Earwigs, Fleas, Ticks, Clover Mites, Brown Dog Ticks, Crickets, Cockroaches, Silverfish, Spiders, Darkling Beetles, and other nuisance insects.

Automatic systems must be programmed so as not to activate and/or release pesticides when people, pets, food or feed are present. A rain sensor and a wind sensor should be installed on the system to control drift and runoff. The system reservoir tank should be locked and a legible copy of the label securely attached to the system reservoir tank, or on the outside of the system timer.

Automatic Spraying System Dilution Table

A System Tank Size (in gallons)	B Conversion Factor (fluid ounce of Concentrate per gallon of water)	A x B = Fluid ounces of Concentrate for that tank size, to be mixed with water to achieve 0.055% pyrethrin solution
1	2.33	2.33
15	2.33	35
20	2.33	46.6
30	2.33	70
55	2.33	128

For best results, set the timer to activate at dusk or dawn when mosquitoes are most active. No more than 3 evening applications recommended.

During the day, manually activate and deactivate the automatic spray system 30 minutes to an hour before outdoor activity is planned.

For high populations of insects, or when treating species that are difficult to control, mix at rates giving higher concentrations (percentage) of pyrethrins up to 0.10% using a conversion factor of 4.08 in table above.

When filling tank, partially fill the system reservoir tank with water, add Concentrate, then fill rest of tank with water and agitate well. System nozzles should deliver fine particle size droplets (aerosol or mist). Nozzles should be rated for a delivery rate not to exceed 1.25 fluid ounces of solution per minute, with one nozzle covering 100 square feet. Set system timer to operate in accordance with equipment directions.

Since high water temperatures can degrade pyrethrins, it is recommended that system tanks be sheltered or insulated from direct exposure to the sun. Use of an EPA approved antimicrobial agent is also recommended. For best performance the spray solution should be in the pH range of 5.5 to 7.0. Avoid setting up system nozzles in a manner that delivers spray solution into swimming pools, or pools containing fish and other organisms that may be sensitive to Pyrethrins.

This product will control insects that are directly contacted by its spray solution. Since many of the targeted insects can migrate in and out of treatment areas, it is recommended that this concentrate be used as part of an Integrated Pest Management (IPM) program. Control practices such as (but not limited to) eliminating breeding and harborage sites, combined with appropriate use of contact insecticides, Insect Growth Regulators (IGRs) and/or residual insecticides will provide the most effective control of target insects.

OUTDOOR USE: follow dilution chart and apply as directed.

0.05% Pyrethrins	add 1 part concentrate to 59 parts water or 2 fluid ounces concentrate / gallon water
0.10% Pyrethrins	add 1 part concentrate to 29 parts water or 4.3 fluid ounces concentrate / gallon water
0.30% Pyrethrins	add 1 part concentrate to 9 parts water or 13 fluid ounces concentrate / gallon water
0.50% Pyrethrins	add 1 part concentrate to 5 parts water or 22 fluid ounces concentrate / gallon water

Caution should be taken to avoid spraying when wind conditions might create a mist that can blow back to applicator. Do not apply this product in or around bodies of water. Spray surfaces and contact as many insects as possible.

SMALL FLYING INSECTS AND OTHER FLYING INSECTS including Flies, Gnats, Mosquitoes, Small Flying Moths, Midges: hold sprayer 18 – 24 inches from shrubs, bushes, grassy areas and other areas infested with these pests. Spray with a slow sweeping motion while moving away from the treated area. To reduce annoyance from these pests, treat open porches and patios in a similar manner. Kills and repels Flies, Mosquitoes including the Culex species that may carry and transmit the West Nile Virus, Midges, and Gnats when used intermittently during outdoor activities. Will keep surrounding areas free from these pests. Allow spray to dissipate before occupying the treated area.

ANTS: Spray ant trails, anthills and other areas of entry and infestation. Spray on ants where possible.

SPIDERS: Spray directly on spiders and webs.

SOWBUGS, MILLIPEDES, BEETLES, EARWIGS, PILLBUGS, CENTIPEDES, SCORPIONS, COCKROACHES, CRICKETS, WATERBUGS, PALMETTO BUGS: Thoroughly spray surfaces and into hiding places such as cracks, crevices, moist areas, openings around pipes, spigots, storage areas, sheds and other areas where these pests tend to congregate.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Store in a cool, dry place. Keep container closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse or equivalent and offer for recycling or reconditioning, or dispose of in a sanitary landfill or by other procedures approved by State and Local authorities.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate.

PHYSICAL OR CHEMICAL HAZARDS

Do not use this product in or on electrical equipment because of shock hazard.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail).

FIRST AID

- IF SWALLOWED:**
- Immediately call a poison control center or doctor.
 - Do not induce vomiting unless told to do so by the poison control center or doctor.
 - Do not give **any** liquid to the person.
 - Do not give anything by mouth to an unconscious person.
- IF ON SKIN**
- Take off contaminated clothing.
- OR CLOTHING:**
- Rinse skin immediately with plenty of water for 15-20 minutes.
 - Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712.



VAMPYRE™ is a trademark of McLaughlin Gormley King Company.
McLaughlin Gormley King Company® is a registered
trademark of McLaughlin Gormley King Company.
Manufactured by McLaughlin Gormley King Company®.
8810 Tenth Avenue North, Minneapolis, MN 55427
037-763